

Heavy Neutral Lepton on Future Muon Collider

Kunfeng Lyu
University of Minnesota

In Collaboration with Peiran Li and Zhen Liu

Muon Collider Physics and Detector Workshop, Fermilab, Dec 15, 2022

Outline

- Neutrino Mass
- Heavy Neutral Leptons
- Search at Future Muon Collider
 - Muon flavor
- Various Experimental Constraints

Origin of Neutrino Mass

- In SM, neutrino is massless. While the experiments have confirmed its tiny mass smaller than $O(0.1)$ eV.

- Effective Operator: Weinberg Operator $\frac{LLHH}{\Lambda}$

- Seesaw mechanism

- Simple Type I
- Inverse seesaw model
- Linear seesaw model

- We choose to work in a simple scenario. Suppose there is heavy neutral lepton. We can parametrize its mass m_N

mixing angle with SM neutrino. $U_l = \sin \theta_l$

$$\mathcal{L} = \mathcal{L}_W + \mathcal{L}_Z + \mathcal{L}_H$$



$$\mathcal{L}_W = \frac{gU_l}{\sqrt{2}} (W_\mu \bar{l}_L \gamma^\mu N + h.c.)$$

$$\mathcal{L}_Z = -\frac{gU_l}{2 \cos \theta_w} Z_\mu (\bar{\nu}_L \gamma^\mu N + \bar{N} \gamma^\mu \nu_L)$$

$$\mathcal{L}_H = -\frac{U_l m_N}{v} h (\bar{\nu}_L N + \bar{N} \nu_L)$$

Our current focus

$$m_N > O(100)\text{GeV}$$



The muon collider can open and probe new region space in the parameter space. even compared to other future colliders!



Search at Muon Collider

- The future muon collider includes 3 TeV and 10 TeV scenarios.
- Clean background, fixed cms energy, excellent environment for the muon flavor HNL
- Here we show the muon-flavor Dirac HNL as benchmark.
- Tools:
 - Using MadGraph 3.4 to simulate and then make analysis
- Effective Vector-Boson Approximation (EVA) or gauge boson PDF has been implemented

Muon Flavor

- Signal: Production of N_μ
Dominated by the t-channel

$$\mu^+ + \mu^- \rightarrow N_\mu + \bar{\nu}_\mu$$



Type	Signal process	$\sigma/U_\mu^2(m_N = 1000\text{GeV})$
t-channel	$\mu^+\mu^- \rightarrow N_\mu\bar{\nu}_\mu$	48.26 pb
VBF	$\mu^+\mu^- \rightarrow \mu^+\mu^-N_\mu\bar{\nu}_\mu$	~ 1 pb
VBF	$\mu^+\mu^- \rightarrow \bar{\nu}_\mu\nu_\mu N_\mu\bar{\nu}_\mu$	~ 0.1 pb

Table 1: Signal for μ flavor at 3 TeV

Decay of N_μ

HNL can promptly decay via neutral current or charged current or to the higgs. Here we select its decay channel to W boson.

$$N_\mu \rightarrow W^+ + \mu^-, W^+ \rightarrow jj$$

We assume the W boson can be well reconstructed from the two jets.

We focus on the final states W^+ and μ^- and reconstruct its invariant mass distribution.

10TeV Background

Dijets can be from either W or Z boson.

Type	Background process	σ/U_μ^2	Cut	Taking account
t-channel	$\mu^+\mu^- \longrightarrow W^+\mu^-\bar{\nu}_\mu$	0.107 pb	default cut	Yes
t-channel	$\mu^+\mu^- \longrightarrow Z\mu^+\mu^-$	0.232 pb	default cuts & missing μ^+	Yes
VBF	$\mu^+\mu^- \longrightarrow \mu^+\mu^-W^+\mu^-\bar{\nu}_\mu$	0.2 pb	default cut ($\mu^+\mu^-$ are missing)	Yes
VBF	$\mu^+\mu^- \longrightarrow \bar{\nu}_\mu\nu_\mu W^+\mu^-\bar{\nu}_\mu$	0.0343 pb	default cut	No


TABLE IV. Background for μ flavor at 10 TeV

Using EVA in MadGraph, especially photon PDF
VBF processes dominates at 10 TeV

Using EVA will lead to t-channel singularity.
So we just generate 2 -> 5 processes directly in MadGraph.

Default cut:
For muon:
PT > 20GeV,
Eta < 2.5

(Real) t-channel singularity




Complex mass
regulation
hep-ph/9509314

The cross section is
proportional to the
beam size.
hep-ph/9601290

We are trying to gain a
better understanding.
(In progress)

(Fake) t-channel singularity: Subtlety of EVA



In 2 to 5 full processes, the intermediate neutrino can never be on-shell.

But in EVA, if we treat the gauge bosons as "real", the neutrino can be on-shell which induces t-channel singularity.

One plausible way is to make the light particles are from the decay of heavy gauge bosons.

Cutflow Analysis

1. We assume the W boson can be well constructed from the dijets (Z boson can fake).
2. Combine the $W^+(Z)$ and μ^- to try to reconstruct the HNL


Impose the mass window:

Require a cut around the heavy neutrino mass.

$$m_N < 500\text{GeV} : \quad m_N \pm 0.1m_N$$

$$m_N \geq 500\text{GeV} : \quad m_N \pm 50\text{GeV}$$

After mass window selection





Optimization cuts I:
Missing transverse
momentum needs to be
less than 500 GeV.

Optimization cuts II:

For the reconstructed W boson.


$1000\text{GeV} < E(W) < 4950\text{GeV}$ which is mass window dependent



Final Results


95% exclusion bounds

10 TeV



Various Bounds

Next step




Short Review of Various Probes

- People have tried to constrain in the U^2 -mN plane via various channels and different machines.
- Cosmo and astrophysical probe: BBN, CMB, etc
- Indirect constraints: branching ratio of SM particles decays, etc
- Direct constraints
 - Production
 - Meson decay, heavy lepton decay
 - (On-shell/Off-shell) Gauge/higgs boson decay
 - Decay
 - Short-lived
 - Long-lived

Existing Bounds

Snowmass Energy Frontier Report: 2211.11084

From Past Experiments



NuTeV: Drell-Yan, NHL decay


CHARM:
Beam-dump: D meson decay,
Wide-band: neutrino beam
colliding with nucleus

DELPHI: Z boson decay

E949: Kaon decay


LHC: Off-shell W/Z decay

Upgraded Bounds




Future Upgraded
Projection on
LHC, NA62 and
DUNE

Various Bounds



Some future
proposed beam-
dump
experiments or
far detector to
probe the long-
lived HNL

Various Bounds



Bounds from the proposed future collider: FCC, CEPC, ILC, LHeC

Conclusion

- Muon Collider is a good platform to probe the TeV scale HNL.
We can open a new region in the parameter space.
- For the muon flavor case, we can probe the $U_{\mu\mu}^2$ down to 10^{-6} for m_N larger than $O(100)$ GeV.
- There are t-channel singularity cases one should be careful to deal with.